

## RECOMMENDATIONS FOR DRY COW THERAPY WITH ANTIBIOTICS

### Definitions

- **Dry cow therapy with antibiotics** means that preferably long-acting antibiotics contained in special udder tubes is infused into all udder quarters of a cow via the teat canal after the last milking during drying-off, i.e. before the start of the dry period.

### Background

- Healthy udders are a prerequisite for sustainable milk production.
- The aim of the recommendations is to ensure that dry cow therapy with antibiotics is used for cows when the treatment result is expected to be good by identifying routines for choosing such cows. A good treatment result will also lead to reduced spread of udder infections in the herd.
- The recommendations also give instructions on how to reduce the risk for contamination and damages at the treatment occasion and give suggestions for routines for follow-up and evaluation of dry cow therapy.
- The routines should always be based on the herd conditions. Thus, the road to the goal (healthy udders in newly calved cows) can vary between herds.

### Recommendations

- Herd specific written routines for dry cow therapy with antibiotics should be produced.
- Dry cow therapy with antibiotics can be considered for cows with subclinical mastitis (elevated SCC without visible signs of mastitis) caused by penicillin-sensitive bacteria.
- Dry cow therapy with antibiotics should be avoided if the time until estimated calving is shorter than 6 weeks at the last milking before the dry period to avoid the risk for a long milk withholding period for cows calving early. The time until estimated calving is especially important in organic herds since the withholding period is doubled in these herds.

■

---

**Contact:** Veterinarian for ruminants, National Veterinary Institute (SVA) 018-67 40 00; Udder health expert, Växa Sverige 010-471 00 00.

**Other:** Published 2021. Updated 2026. The project was financed by the National Board of Agriculture via SvarmPat.

- Cows considered for dry cow therapy with antibiotics are selected based on udder health class (juverhälsoklass; JHKL) at the last test milking before drying off
  - o If information on JHKL is not available, the selection can be based on an average of 2 to 3 evenly distributed on-line cow cell count measurements during 2-3 months before drying-off. See Table A for translation of cow somatic cell counts to JHKL.
  - o If the herd has no access to JHKL or cow somatic cell counts the selection can be based on testing with California Mastitis Test (CMT) according to Table B (see description and video here: <http://www.juverportalen.se/media/1136/cmt-20190412.pdf>). Knowledge on the best way to select cows using CMT is not available. A suggestion is to examine the cows approximately 2 weeks before drying-off and at the last milking before the start of the drying-off. If the cow has at least one udder quarter with CMT 2–5 at both occasions the cow may be considered for treatment.
- Perform the dry cow therapy directly after the end of the milking session to reduce the risk of contamination of milk and facilitate correct performance of the treatment.
- Practical performance of dry cow therapy with antibiotics:
  - o All 4 udder quarters are treated after the last milking after careful cleaning of the teat ends according to the following routine:
    - It is preferable that two persons are present, and that one person assists the other with cotton wool and tubes.
    - Wash hands before treatment and use new disposable gloves for each cow.
    - Remove visible dirt from the teats with dry or moist paper when needed.
    - Dip the teats in teat disinfectant and leave on for 30 seconds. Wipe teats with an individual paper towel per teat. Start with the teat furthest away and end with the closest teat.
    - Prepare one teat at the time according to the following description. Start with the teat closest to you.
      - Wipe the teat end carefully, especially around the opening of the teat canal, with several pieces of cotton wool moistened with 70% alcohol until no dirt is visible on the cotton wool.
      - Let the teat dry for 30 seconds.
    - Perform the treatment of each teat/udder quarter as follows:
      - Remove the top part of the cap on the tube so that the end of the tube tip is exposed. Avoid contaminating the tip by touching the udder skin or fingers.
      - Hold the teat with one hand and angle it so it is easy to see the opening of the teat canal.
      - Carefully insert the end of the tube tip into the teat canal.
      - Infuse all the contents of the tube into the teat.
      - Carefully withdraw the tip of the tube from the teat canal.
      - Gently massage the teat and the lower part of the udder quarter upwards to facilitate spread of the product within the udder quarter.
    - Teat dip/spray all teats with teat disinfectant directly after treatment.

**Contact:** Veterinarian for ruminants, National Veterinary Institute (SVA) 018-67 40 00; Udder health expert, Växa Sverige 010-471 00 00.

**Other:** Published 2021. Updated 2026. The project was financed by the National Board of Agriculture via SvärmPat.

- Place the cow in a suitable group decided by her udder health status during the dry period and the time around calving.
- Control the udder and the general condition of the cow and apply teat dip/spray twice daily for 14 days according to the routine for drying-off (see separate document).
- All cows that had dry cow therapy with antibiotics should be checked after calving as follows:
  - o Perform CMT day 3–4 after calving (see instruction and video here:  
<http://www.juverportalen.se/media/1136/cmt-20190412.pdf>)
    - If the CMT-reaction is 2–5 milk samples for bacteriological examination are taken from such quarters. The cow should be milked after healthy cows.
  - o Control the cow somatic cell counts at the first test milking
    - If the cow somatic cell count is above 150 000 cells/ml the udder is examined using CMT and milk samples are taken as described above. The cow should be milked after healthy cows.
- Evaluate the herd routines by regular follow-up of the udder health after calving (e.g. the proportion of cows with CMT 2–5 day 3–4 after calving; the proportion of cows with clinical mastitis during the first month; differences in somatic cell counts at the first test milking compared to the one before drying-off (e.g. the proportion of new infections or the proportion of cured (cut-off healthy/sick 150 000 cells/ml); results from bacteriological examinations).

*Table B. Basis for selection of cows considered for treatment with dry cow antibiotics or internal teat sealants based on udder quarter California Mastitis Test (CMT) approximately 2 weeks before drying-off and at the last milking before drying-off*

CMT	Interpretation	Cell count x 1000 cells/ml	Dry cow antibiotics	Teat sealants
1	Negative	<200	NO	YES*
2	Trace	150–500	YES** Depending on culture findings and cell count history	NO***
3	Weak positive	400–1500		
4	Distinct positive	800–5000		
5	Strong positive	>5000		

Schalm, Carroll and Jain, 1971; with addition of Scandinavian scoring (1–5). \* If all udder quarters of a cow have CMT 1 and the herd has problems with environmental udder infections. \*\* If the cow has at least one udder quarter with CMT 2–5 at both samplings the cow may be considered for treatment. \*\*\* In herds with environmental udder infections a combination of dry cow antibiotics and internal teat sealants may sometimes be considered.

**Contact:** Veterinarian for ruminants, National Veterinary Institute (SVA) 018-67 40 00; Udder health expert, Växa Sverige 010-471 00 00.

**Other:** Published 2021. Updated 2026. The project was financed by the National Board of Agriculture via SvarmPat.